



UNIVERSITY
of HAWAII®

MĀNOA

**University of Hawai`i at
Mānoa Department of
Economics
Working Paper Series**

Saunders Hall 542, 2424 Maile Way,
Honolulu, HI 96822
Phone: (808) 956 -8496
www.economics.hawaii.edu

Working Paper No. 16-15

Health and Health Inequality during the Great
Recession: Evidence from the PSID

By
Huixia Wang
Chenggang Wang
Timothy J. Halliday

September 2016

Health and Health Inequality during the Great Recession: Evidence from the PSID

Huixia Wang

Hunan University, School of Economy and Trade

Chenggang Wang

University of Hawaii at Manoa, Department of Economics

Timothy J. Halliday

University of Hawaii at Manoa, Department of Economics

University of Hawaii Economic Research Organization

IZA

First Version: July 6, 2016

Last Update: September 8, 2016

Abstract

We employ granular information on local macro-economic conditions from the Panel Study of Income Dynamics to estimate the impact of the Great Recession on health and health-related behaviors. Among working-aged adults, a one percentage point increase in the county-level unemployment rate resulted in a 2.4-3.2% increase in chronic drinking, a 1.8-1.9% decrease in mental health status, and a 7.8-8.9% increase in reports of poor health. Notably, there was heterogeneity in the impact of the recession across socioeconomic groups. Particularly, obesity and overweight rates increased for blacks and high school educated people, while there is weak evidence that they decreased for whites and the college educated. Along some dimensions, the Great Recession may have widened some socioeconomic health disparities in the United States.

Keywords: Great Recession, Health Behaviors, Health Outcomes, Obesity, Inequality

I. Introduction

Recessions are a major source of systematic risk to households. Because they affect large groups of people at once, they are very difficult to insure. Moreover, due to moral hazard problems, public insurance schemes like unemployment insurance only provide limited recourse to the unemployed. As a consequence, recessions can have serious, adverse impacts on household and individual welfare.

One of the more commonly studied of these potential impacts is the effect of recessions on human health. Early work on the topic indicated that poor macroeconomic conditions raised mortality rates substantially (*e.g.* Brenner 1979). However, seminal work by Ruhm (2000) pointed out severe methodological shortcomings in this earlier work and he showed that, once these issues are corrected, mortality rates tend to *decline* during recessions so that mortality rates are actually pro-cyclical in the aggregate data.¹ Improved health-related behaviors due to relaxed time constraints and tightened budget constraints was cited as a mechanism driving these results by Ruhm (2000, 2005), although subsequent work by Stevens, *et al.* (2015) suggested that higher rates of vehicular accidents and poor nursing home staffing during robust economic times were the primary mechanisms. Notably, more recent work by Ruhm (2015) has shown that mortality rates for many causes of death did not decline during the Great Recession and that mortality due to accidental poisoning actually increased.² Importantly, all of these studies utilize aggregate state-level mortality and unemployment rates and so their unit of analysis is a state/time observation.

On the other hand, studies that are based on individual-level data tend to show that health and health-related behaviors worsen during recessions. For example, Gerdtham and Johannesson (2003, 2005) use micro-data and show that mortality risks increase during recessions for working-aged men. Similar evidence over the period 1984-1993 is provided for the United States by Halliday (2014) who used the Panel Study of Income Dynamics (PSID). In a similar

¹ This result has been replicated in other countries such as Canada (Ariizumi and Schirle 2012), France (Buchmueller, *et al.* 2007), OECD countries (Gerdtham and Ruhm 2006), Spain (Tapia Granados 2005), Germany (Neumayer 2004), and Mexico (Gonzalez and Quast 2011).

² Related to this finding, Bassols, *et al.* (2016) showed that the Great Recession increased legal and illegal drug use in Spain.

vein to these studies, Jensen and Richter (2003) showed that pensioners who were adversely affected by a large-scale macroeconomic crisis in Russia in 1996 were 5% more likely to die within two years of the crisis. Related, Charles and DeCicca (2008) use the National Health Interview Survey (NHIS) and MSA-level unemployment rates and show that increases in the unemployment rate were accompanied by worse mental health and increases in obesity. Hence, while the macro-based studies tend to be somewhat conflicted, the micro-based studies indicate that the uninsured risks posed by recessions have real, adverse impacts on human health.

In this study, we investigate the impact of the Great Recession of 2007-2009 on health and health-related behaviors using the PSID. This is an important episode to study since this recession was the deepest and longest recession during the post-war period. In fact, Farber (2015) estimates that, over this period, one in six workers lost their job at least once. From trough to peak, the unemployment rate increased from 4.6 percent to 9.3 percent which is the largest increase during the post-war period. To illustrate, we present Figure 1 which shows the unemployment rate during this period. This figure clearly indicates that the recession of 2007-2009 was the most severe. In addition, as shown in Figure 2, unemployment duration during the most recent recession was also, by far, the longest of any recession since World War II peaking at just over 40 weeks.

In this paper, we build on recent work by Tekin, *et al.* (2013) who also study the health impact of the Great Recession. We build on their work by using county-specific unemployment rates as opposed to state-specific rates. Interestingly, we find larger and more significant results for similar outcomes. One explanation for this is that state-specific unemployment rates could be viewed as error-ridden proxies for county-specific unemployment rates and so the estimates in the Tekin, *et al.* (2013) study might suffer from attenuation biases. In addition, we use the PSID's geocode file which allows us to include county-specific fixed effects which control for a rich set of confounding variables at a more granular level. Importantly, this is the first study that looks at the impact of the Great Recession on individual health using county-level information on macroeconomic conditions.

In this sense, our study is also related to Charles and DeCicca (2008) who employ MSA-specific

fixed effects and individual-level data from the NHIS. However, there are important differences between our study and theirs. First, their study does not consider the Great Recession and as indicated by Ruhm (2015), this most recent recession may have impacted human health in considerably different ways than previous recessions. Second, because we use a panel, we are also able to employ individual fixed effects in robustness checks. Third, they limit their study to the 58 largest MSA's in the United States, whereas we use a sample that is more representative of the United States. Finally, while we do replicate their key findings that mental health and obesity worsen during recessions, we also provide evidence that drinking and smoking behaviors were affected by the Great Recession, albeit in complicated ways.

In particular, we document that the Great Recession had more adverse consequences on health-related behaviors for blacks and less educated people than it had on whites and more educated people. Obesity, smoking, and drinking all increased for blacks. In contrast, the effects of the recession on drinking were smaller for whites. Moreover, there is weak evidence that the recession actually *reduced* obesity and smoking prevalence for whites. In a similar vein, we show that the Great Recession had larger adverse effects on drinking, obesity, and mental health for less educated people. Interestingly, the fact that the recession appeared to reduce obesity rates for more privileged groups but increased them for the less privileged is consistent with recent findings by Coleman and Dave (2013). Specifically, they show that recessions increase physical exertion for people who are more likely to be employed in white collar professions but decrease it for people who more likely to be employed in blue collar professions. Consequently, another impact of the Great Recession was that, along many dimensions, it widened socioeconomic disparities in health-related behaviors.

The balance of this paper is organized as follows. In the next section, we discuss our data. After that, we describe our empirical methods. We then present our findings. Finally, we conclude.

II. Data

We utilize data from the PSID which is a national longitudinal study that collects individual-specific information on health, demographic, and socioeconomic outcomes that is run by the

University of Michigan. The PSID began in 1968 with interviews of about 5000 families and has continued to interview their descendants since then. To obtain county-specific information, we use the county identifier file from the PSID.³ We utilize the 2003, 2005, 2007, 2009, 2011 and 2013 waves. We employ these waves because the 2007 and 2009 waves contain the recession and we have two waves prior to the recession (2003 and 2005) and two waves after the recession (2011 and 2013). Because only heads of household and their spouses were asked the health-related questions in the survey, we limit our sample to household heads and their spouses. We employ county-level unemployment rates from the Local Area Unemployment Statistics (LAUS) of the Bureau of Labor Statistics (BLS) which were then merged into the PSID for each year using PSID's geocode file.

For most of the estimations, we restrict the sample to people with strong labor force attachments which we define to be people between ages 25 and 55. In addition, we further restrict this sample by dropping retired and disabled people, students, and housewives. We also present some estimations for people age 65 or older. The idea for using this sample is that this sub-sample has weaker labor force attachments and so if the impact of the recession on health is operating through the labor market then we should see attenuated effects in this population. Because the goal of this exercise is to see if the recession impacted people with weak labor force attachments, we included the retired, disabled, students (to the extent that there are full-time students older than 65), and housewives.

Descriptive statistics for our sample are reported in Table 1. The data can be broadly categorized under the following rubrics: health-related behaviors, health outcomes (physical and mental), and demographics. The demographic variables are fairly self-explanatory and are listed in the bottom portion of the table. The health variables require a bit more explanation which we provide in the next two subsections. With a few exceptions, the outcomes that we use closely follow Tekin, *et al.* (2013).

The county-level unemployment rate was obtained from the LAUS of the Bureau of Labor

³ See <http://simba.isr.umich.edu/restricted/ProcessReq.aspx> for details.

Statistics (BLS). We collected 3,218 county unemployment rates from every other year between the years of 2003 to 2013 which corresponds to the years of our PSID sample. In our sample, the average county-level unemployment rate was 6.95 percent with a standard deviation of 2.75 percent indicating that there is substantial variation in county-level unemployment rates in our data. Moreover, a regression of the county-level unemployment rate onto county fixed effects has an R^2 of 47.22% indicating that over half of the variation of the county-level unemployment rate is within counties which is critical for our research design.

Health-related Behaviors

The PSID includes numerous questions about alcohol and smoking consumption. Specifically, respondents were asked how often they drank and individuals who reported drinking several times a week or everyday were categorized as a “chronic drinker.” The PSID survey also asked respondents about the typical number of drinks consumed per day. Following standard guidelines, we define men who drink three to four drinks or more per day and women who consume one to two drinks or more per day as “heavy drinkers.”⁴ A “current drinker” is defined as an individual who reported having had a drink in the last year. Similarly, we define a “current smoker” as someone who has ever smoked during the last year. Since the definition of light or heavy smoker varies widely, we define a “heavy smoker” as individuals who smoke six or more cigarettes per day.⁵

In addition, we consider physical activity. The PSID provides information about vigorous physical activity (which includes activities such as heavy housework, aerobics, running, swimming, and bicycling), light physical activity (which includes activities such as walking, dancing, gardening, golfing, and bowling), and strengthening activities (which includes activities that are specifically designed to strengthen muscles such as lifting weights). Each of these three variables is binary and is set to unity if the respondent reported any of these types of activities.

⁴ According to the definition from the Centers for Disease Control and Prevention, heavy drinking is defined as consumption of 15 or more drinks per week for men and consumption of 8 or more drinks per week for women. See <http://www.cdc.gov/alcohol/faqs.htm#heavyDrinking>.

⁵ Husten (2009) documents that the definition of light smoking ranged from "smoking 1-39 cigarettes per week" to "smoke 10-20 cigarettes per day."

Finally, we use weekly hours of housework as an additional measure of physical activity.

Health Outcomes

We use the *K6 Non-specific Psychological Distress* scale which was also used by Charles and DeCicca (2008). Recent work by Tekin, *et al.* (2013) also looks at mental health as an outcome, although they use a different depression index. This K6 distress scale includes six questions designed to measure different markers of psychological distress including reports of feelings of effortlessness, hopelessness, restlessness, sadness, and worthlessness during the past 30 days. The K6 distress scale is a weighted sum of these six outcomes. Kessler, *et al.* (2003) has shown that the K6 scale is at least as effective as a number of other depression scales in predicting serious mental health problems.

The physical health outcomes that we employ include variables based on self-reported health status (SRHS) and body mass index (BMI). SRHS is a categorical variable that takes on integer values between one and five where one is excellent and five is poor. We transform the SRHS variable into two binary indicator variables for excellent (SRHS equal to one or two) and poor (SRHS equal to four or five) health. The BMI variable is transformed into three dummy variables for being overweight ($BMI \geq 25$), obese ($BMI \geq 30$), and severely obese ($BMI \geq 35$).

III. Methodology

To estimate the effect of the Great Recession on health outcomes and health-related behaviors, we employ a linear probability model. If we let i denote the individual, c the county, s the state, and y the year, the basic estimation model is:

$$H_{icsy} = \beta_0 + \beta_1 U_{cy} + \beta_2 X_{iy} + \delta_c + \delta_y + \delta_s * t + \varepsilon_{icy}. \quad (1)$$

The dependent variable, H_{icsy} , is a health outcome or behavior. The county-specific unemployment rate in a given year is U_{cy} . The vector, X_{iy} , contains individual-specific control variables including age, gender, race, marriage status, education, and labor income. We also

include county and year dummies which are denoted by δ_c and δ_y . Finally, we include state-specific time trends which are denoted by $\delta_s * t$. We estimate two different specifications of equation (1) both with and without the state-specific trends which has the advantage of controlling for potentially confounding within state trends but the disadvantage of eliminating potentially meaningful exogenous variation in the county-level unemployment rates. Finally, we employ the weights provided by the PSID when estimating these models.

We compute the standard errors using two-way clustering by year and county. When we cluster by county, this accounts for any potential serial correlation within counties or individuals. When we cluster by year, this accounts for spatial correlations within a year. Note that this is a very conservative approach to computing the standard errors.

Our study like Charles and DeCicca (2008) does a comprehensive job of controlling for heterogeneity across local labor markets. Importantly, Tekin, *et al.* (2013) and Ruhm (2005), which also closely align with our own study, only control for state fixed effects which only accounts for the state-level and time-invariant confounders. Clearly, the use of state fixed effects may be too coarse since potential confounders such as education and health infrastructure, culture, demographic composition, and weather may vary at a finer geographical level. For example, Asians are about one third of the population in San Francisco whereas they are only 0.4% of the population of Sierra County in California. Another example is that within states, particularly in the South, some counties are “dry” which means that alcohol cannot be purchased within them. Given that many of our outcomes relate to alcohol consumptions, this is another important potential county-specific confound. Simple inclusion of state fixed effects would not account for these within state confounders.

We also adopt a more comprehensive approach to addressing heterogeneity by including individual fixed effects which subsume the county fixed effects. This approach has the advantage of controlling for a greater amount of unobserved confounding variables than the county fixed effects. However, it comes with the cost of wasting important and exogenous variation in the data as has been argued by Deaton (1997) and Angrist and Pischke (2008). Importantly, even if the individual fixed effect is uncorrelated with the right-hand side variables

of interest, lack of variation within individuals in the dependent variable will bias individual fixed effects estimates towards zero. As such, we view the results with the individual fixed effects as a robustness check for our core results and we primarily focus on the results with the county fixed effects for most of the paper.

IV. Results

Core Results

Our core estimation results are reported in Table 2. In the first column, we include county and year fixed effects. In the second column, we further include state-specific trends. Each cell corresponds to a separate estimate of our parameter of interest, β_1 , and each row corresponds to a separate outcome.

First, we see strong evidence that increases in the unemployment rate during this period increased drinking and, particularly, problematic drinking. In the first column, we see that increases in unemployment are positively associated with increased prevalence of drinking and chronic drinking. In the second column, we see that it is associated with increases in all three types of drinking behavior that we consider. Particularly, a one PP increase in the unemployment rate is associated with a 0.6-0.8 PP increase in chronic drinking and a 0.5-0.6 PP increase in current drinking. This constitutes a 2.4-3.2% increase in the prevalence of chronic drinking and a 0.71-0.86% increase in current drinking.

Next, we do not see strong evidence that increases in the unemployment rate impact smoking behaviors or reports of physical activities. We do see evidence that heavy smoking *decreased* in the both specifications, but only the estimate without the state specific trends is significant. However, there is no evidence that the recession impacted reports of currently being a smoker. Finally, we look at four different measurements of physical activity (vigorous physical activity, light physical activity, strengthening activity, and housework hours) using our two different specifications and we see only one estimate for strengthening activity that is negative and significant at the 10% level. On the whole, there is little to no evidence that the recession impacted physical activities or smoking.

Next, we do see some evidence that there was a deterioration of mental health outcomes induced by the recession. Consistent with Teft (2011), when we look at the K6 index, we see strong evidence that the recession led to worsened mental health outcomes using both specifications. The estimates indicate that a one PP increase in the unemployment rate resulted in an increase in the K6 index of 1.8-1.9%. This apparent deterioration in mental health is consistent with the observed increase in drinking that we just saw which may have been a way of coping with the stress induced by the recession.

In addition, the recession did have strong effects on self-reported health status using both specifications. We see that a one PP increase in the unemployment rate is associated a 0.9 PP decrease in self-reports of excellent health and a 0.7-0.8 PP increase in self-reports of poor health. This constitutes a 1.5% increase in the probability of reporting good health and a 7.8-8.9% increase in the probability of reporting poor health. This result is consistent with Halliday (forthcoming) who showed that fluctuations in earnings have causal impacts of self-rated health.

Finally, there is no evidence that the recession impacted obesity in this table. None of our three measures of obesity and overweight were impacted. However, it is important to note that this could mask important variation across demographic subgroups as we will see.

In Table 3, we estimate the same models that we estimated in the previous table for the subsample that is 65 or older. This population is, by-and-large, retired and so has weak labor force attachments. Looking at the results from our preferred specification in the second column, we see that none of the estimates are significant at the 5% or 1% levels and only the estimate of the coefficient on severe obesity is significant at the 10% level. In the first column where we exclude the state-specific time trends, we see that the coefficient estimates on vigorous physical and strengthening activities are positive and significant but only at the 10% level and this result is not robust to the inclusion of the state-specific trends. On a similar note, there is evidence in the first column that the recession improved self-rated health but, once again, this finding is not robust to the inclusion of state-specific time trends. On the whole, the results in this table do not indicate that the Great Recession had systematic, negative effects on an elderly population with

weak labor force attachments. This seems to indicate that the effects of the recession on health ran primarily through the labor market.

Next, in Table 4, we conduct a series of robustness checks. The table contains four sets of estimations. The first two sets include individual fixed effects both with and without state-specific trends. The second two are estimates of county fixed effects models using a sub-sample of people who did not change counties while in the sample, again, both with and without state-specific trends. These two exercises address a concern discussed in Halliday (2007) in which healthier people migrate in response to recessions. So, if the estimate indicates that the recession is adversely impacting health, then the bias stemming from selective out-migration would make it appear as if the recession is worse for human health than it actually is. On the other hand, the reverse is true if the estimates indicate that some aspect of health or health behavior improves during recessions. Note that, while these exercises do address this concern, they also place much heavier demands on the data which, most likely, will result in higher standard errors and, hence, reduced power.

The results in Table 4, on the whole, buttress the results in Table 2. First, we still see strong evidence that problematic drinking increased during the recession when looking at the estimates using the non-mover sample and we see similar (albeit somewhat weaker) evidence when employing individual fixed effects in the first two columns. Interestingly, the results in this table suggest that smoking and obesity actually decreased during the recession. Particularly, a one PP increase in the unemployment rate reduced the prevalence of heavy smoking by between 0.3 and 0.5 PP and obesity by 0.5 PP. Note that we did not see significant effects on smoking or obesity in Table 2. This is consistent with our discussion of the selection bias induced by migration in the previous section. Healthier people leave depressed areas which will create a positive bias in the smoking and obesity estimates in Table 2. Finally, we also see negative effects of the recession on SRHS and the K6 index, although the effects on the K6 index are not significant in the individual fixed effects specifications.

Results by Demographic Group

We begin this sub-section by estimating equation (1) using a dummy variable for currently being employed for our sample of 25-55 year olds. We estimate the model by race (black and white), gender, and education (people with 12 or fewer years of schooling and people with college degrees).⁶ The point of this exercise is to obtain an idea of which groups were most impacted by the recession and to compare this to the health impacts of the recession by demographic sub-groups. Not surprisingly, all six of our estimations delivered coefficient estimates on the county-level unemployment rate that were significant at 1% level.

The point-estimates are reported in Figure 3. We see that a one PP increase in the unemployment rate reduced the probability of being employed in the PSID by 0.6 PP for whites and 0.9 PP for blacks. Similarly, we see that the corresponding marginal effects for men and women are 0.6 and 0.8 and the estimates by education are 0.6 and 0.7 for the college and high school educated. The overall pattern is that, while the recession had a broad impact, women and more disadvantaged socio-economic groups were the most impacted by it.

In Table 5, we report our estimation results of both specifications from Table 2 separated by race. First, looking at the drinking outcomes, we see that the recession had much stronger effects on the drinking behavior of black people than on white people. For example, when we included state-specific trends in the second and fourth columns, we see that a one PP increase in the unemployment rate is associated with a 0.9 PP increase in chronic drinking for whites and a 2.2 PP increase for blacks. Similarly, in the same specifications, the probability of being a current drinker increased by 0.5 PP for whites and 1.8 PP for blacks. We also see that a one PP increase in the unemployment rate resulted in a 1.1-1.7 PP increase in the prevalence of heavy drinking for blacks but had no effects on whites. We also see that blacks report a 1.1-1.3 PP increase in the probability of being a current smoker in response to a one PP increase in the unemployment rate. Interestingly, there is weak evidence that whites smoked *less* in response to the recession; in the first column, we see that the probability of being a heavy or a current smoker decreased by 0.4-0.6 PP. As before, we see no evidence, at the 5% or 1% level, that the recession impacted any of our physical activity measures. Turning to the mental health outcomes, we see that the recession had large effects on whites but no effects on blacks. In a similar vein, while there is

⁶ We estimated the more parsimonious specification that excluded the state-specific time trends.

some evidence that blacks reported worse self-rated health in column three, the bulk of the effects on SRHS appears to have been for the white sub-population. Specifically, we see that a one PP increase in the unemployment rate is associated with a 0.9 PP decrease in the probability of reporting excellent health and a 0.7-0.8 PP increase in the probability of reporting poor health for whites.

Finally, as with smoking, we see an interesting contrast in how the recession impacted obesity rates by race. Looking at whites in the first column, there is evidence that the rate of obesity and severe obesity *declined* by 0.5 and 0.3 PP, respectively. However, this result is not robust to the inclusion of state-specific trends. In contrast, all three measures of obesity increased in response to increases in the unemployment rate for blacks. This is true in both specifications for columns three and four. These effects vary in a narrow band between 1.1 and 1.5 PP.

In Table 6, we present estimates by gender. First, the recession impacted drinking for both men and women, although in different ways. For men, the bulk of the impact was on moderate drinking. Indeed, we see that there is weak evidence that chronic drinking increased by 0.5-0.8 PP and stronger evidence that current drinking increased by 0.7 PP for men. In contrast, for women, the probabilities of being a current or heavy drinker were unaffected, but the probability of being a chronic drinker increased by 0.8 PP and this estimate is significant at the 95% level in both specifications. Next, there is little evidence that the recession impacted smoking or physical activity for either gender. There is some evidence that the recession impacted mental health outcomes, but looking at the point-estimates of the coefficient on the K6 index, we do not see any discernible gender differences as the point-estimates are all similar. In the same spirit, we see that fluctuations in the unemployment rate impacted SRHS for both genders, but the effects were larger for women. Specifically, we see that a one PP increase in the unemployment rate is associated with a 0.7-1.0 PP increase in the probability of reporting being in poor health for women and 0.4-0.5 PP for men. Similarly, we see a decrease in the likelihood of excellent health of 0.9 PP for women and 0.7 PP for men. Finally, we see no evidence that the recession impacted obesity rates for men, but we do see some evidence that the recession reduced obesity rates for women as the probability of being obese declined by between 0.7-0.8 PP for women.

Finally, we investigate differential impacts of the Great Recession by education levels in Table 7. Specifically, we estimate the models separately for people with college degrees and people with at most a high school degree. First, we see that the bulk of the effects of the recession on drinking occurred for the least educated. We see no effects for people with college degrees but we see that there was a 0.6-0.8 PP increase in the probability of being a current drinker and a 0.6-1.0 PP increase in the probability of being a chronic drinker for people with at most a high school degree. As before, we see little to no evidence that the recession impacted either smoking or physical activity for either educational group. Next, we see that the bulk of the effects of the recession on mental health outcomes occurred for the least educated as evidenced by the estimates of the coefficient on the K6 index; the point estimates in the third and fourth columns for the high school educated group are 0.064 and 0.53, respectively, whereas they are 0.018 and 0.040 in the first two columns for the college-educated group. A similar pattern exists for SRHS. We see no effects for the college-educated, but we do see a 1.0-1.2 PP reduction in reports of excellent health and a 0.7-0.8 PP increase in reports of poor health for the least educated. Finally, and similar to our results broken down by race, we see that the recession reduced the probability of being overweight for the college educated by 1.1 PP in the first specification but increased the likelihood of being overweight for people with at most a high school degree by 0.6 PP in both specifications.

These results underscore some of the distributional impacts of the Great Recession in ways that have not hitherto been articulated in the literature. For many important health outcomes, the recession appears to have exacerbated health disparities between blacks and whites and across educational groups. By race, we saw that the recession had larger impacts on drinking, smoking, and obesity rates for blacks than for whites. Moreover, there is some evidence that the recession actually resulted in lower obesity rates for whites and women. However, the recession impacted subjective health measures (*i.e.* the K6 index and the SRHS measures) more for whites than blacks. In addition, the results by educational group indicate that there were larger effects on drinking, mental health, SRHS, and obesity for the least educated. The results for subjective health outcomes by race notwithstanding, these results are consistent with Figure 3 in that blacks and high school educated people were most impacted by the recessions and these groups drank more and experienced a greater deterioration in their health status in response to the exigencies

posed by the Great Recession.

V. Conclusions

In this paper, we showed that, overall, the Great Recession resulted in worse health outcomes. We built on previous work by employing more granular information on local macroeconomic conditions by using the geocode file from the Panel Study of Income Dynamics. For a population of working-aged adults, we showed that a one percentage point increase in the county-level unemployment rate resulted in a 2.4-3.2% increase in chronic drinking, a 1.8-1.9% increase in the K6 depression index, and a 7.8-8.9% increase in the probability of reporting poor health. Importantly, however, these effects mask that there was considerable heterogeneity in the recession's effects across socioeconomic groups.

As discussed above, it is well documented that aggregate mortality rates are pro-cyclical (*e.g.* Ruhm 2000, Stevens, *et al.* 2015). Ruhm (2000) suggested that one of the mechanisms behind this counter-intuitive finding was that health-related behaviors improved during recessions since relaxed time constraints would enable more time spent exercising and tightened budget constraints would result in less money spent on vices such as alcohol and cigarettes. Indeed, Ruhm (2005) provided evidence that tobacco consumption decreases and physical activity increases during economic lulls in the BRFSS over the period 1987-2000, although Tekin, *et al.* (2013) show that this relationship has severely weakened or become zero over the period 2005-2011 in the BRFSS. Our results, on the whole, are not consistent with these findings in that we find that SRHS, mental health, and problematic drinking behavior all worsen during recessions for people of prime working age, although some of our findings indicate that some health-related behaviors may have improved for whites and the college-educated. It is also important to note that more recent work by Ruhm (2015) has shown that mortality rates have not been pro-cyclical since the early 2000's. In this sense, since we look at the period 2003-2013, our work need not be viewed as being at loggerheads with Ruhm (2000).

While this work may seem to be at odds with many of the studies that rely on aggregate data

(*e.g.* state-level mortality rates) discussed above, the results are consistent with studies on the impact of recessions on mortality that use micro-data. For example, Gerdtham and Johannesson (2003, 2005) use individual-level Swedish administrative data and find that mortality risk increases during economic downturns for working-aged men. Similarly, Halliday (2014) uses the PSID and finds that a one PP increase in the unemployment rate results in a 6% increase in one-year mortality risk at baseline also for working aged men.

Our findings on the effects of the Great Recession on obesity are also consistent with other findings in the literature. For example, and similar to our own findings, Charles and DeCicca (2008) find the obesity tends to increase during recessions for men with low *ex ante* employment probabilities and for black men. Evidence for a mechanism underlying this finding is provided by Coleman and Dave (2013) who show that, while exercise hours increase during recessions, for less educated individuals, this does not compensate for total loss in physical exertion due to the loss of physically demanding jobs.

Related to this, an important conclusion of our work is that different socioeconomic groups responded to the Great Recession in very different ways. As already discussed, the Great Recession increased obesity rates for black and high school educated people but there is weak evidence that it decreased them for white and college educated people. Similar results also obtain for smoking by race; whites smoked less in response to the recession, whereas blacks smoked more. Finally, the recession adversely impacted the drinking behaviors of black and white people and of less educated and more educated people; however, the magnitudes were much larger for black and less educated people. In this sense, the Great Recession appears to have exacerbated many socio-economic health disparities in the United States.

References

Angrist, Joshua D., and Jörn-Steffen Pischke. *Mostly harmless econometrics: An empiricist's companion*. Princeton university press, 2008.

Ariizumi, Hideki, and Tammy Schirle. "Are recessions really good for your health? Evidence from Canada." *Social Science & Medicine* 74, no. 8 (2012): 1224-1231.

Bassols, Nicolau Martin, and Judit Vall Castelló. "Effects of the Great Recession on drugs consumption in Spain." *Economics & Human Biology* 22 (2016): 103-116.

Brenner, M. Harvey. "Mortality and the national economy: A review, and the experience of England and Wales, 1936-76." *The Lancet* 314, no. 8142 (1979): 568-573.

Buchmueller, Thomas C., Michel Grignon, Florence Jusot, and Marc Perronnin. *Unemployment and mortality in France, 1982-2002*. Centre for Health Economics and Policy Analysis, McMaster University, 2007.

Charles, Kerwin Kofi, and Philip DeCicca. "Local labor market fluctuations and health: is there a connection and for whom?." *Journal of Health Economics* 27, no. 6 (2008): 1532-1550.

Coleman, Gregory, and Dhaval Dave. "Exercise, physical activity, and exertion over the business cycle." *Social Science & Medicine* 93 (2013): 11-20.

Deaton, Angus. *The analysis of household surveys: a microeconomic approach to development policy*. World Bank Publications, 1997.

Farber, Henry S. *Job loss in the Great Recession and its aftermath: US evidence from the displaced workers survey*. No. w21216. National Bureau of Economic Research, 2015.

Gerdtham, Ulf-G., and Magnus Johannesson. "A note on the effect of unemployment on mortality." *Journal of health economics* 22, no. 3 (2003): 505-518.

Gerdtham, Ulf-G., and Magnus Johannesson. "Business cycles and mortality: results from Swedish microdata." *Social science & medicine* 60, no. 1 (2005): 205-218.

Gerdtham, Ulf-G., and Christopher J. Ruhm. "Deaths rise in good economic times: evidence from the OECD." *Economics & Human Biology* 4, no. 3 (2006): 298-316.

Gonzalez, Fidel, and Troy Quast. "Macroeconomic changes and mortality in Mexico." *Empirical Economics* 40, no. 2 (2011): 305-319.

Granados, José A. Tapia. "Recessions and mortality in Spain, 1980–1997." *European Journal of Population/Revue européenne de Démographie* 21, no. 4 (2005): 393-422.

Halliday, Timothy J. "Business cycles, migration and health." *Social Science & Medicine* 64, no. 7 (2007): 1420-1424.

Halliday, Timothy J. "Unemployment and Mortality: Evidence from the PSID." *Social Science & Medicine* 113 (2014): 15-22.

Halliday, Timothy J. "Earnings Growth and Movements in Self-Reported Health," forthcoming *Review of Income and Wealth*.

Husten, Corinne G. "How should we define light or intermittent smoking? Does it matter?." *Nicotine & Tobacco Research* 11, no. 2 (2009): 111-121.

Jensen, Robert T., and Kaspar Richter. "The health implications of social security failure: evidence from the Russian pension crisis." *Journal of Public Economics* 88, no. 1 (2004): 209-236.

Kessler, Ronald C., Gavin Andrews, Lisa J. Colpe, Eva Hiripi, Daniel K. Mroczek, S-LT Normand, Ellen E. Walters, and Alan M. Zaslavsky. "Short screening scales to monitor population prevalences and trends in non-specific psychological distress." *Psychological medicine* 32, no. 06 (2002): 959-976.

Neumayer, Eric. "Recessions lower (some) mortality rates: evidence from Germany." *Social Science & Medicine* 58, no. 6 (2004): 1037-1047.

Ruhm, C. J. "Are Recessions Good for Your Health?" *The Quarterly journal of economics*, 115, no. 2 (2000): 617-650

Ruhm, Christopher J. "Healthy living in hard times." *Journal of Health Economics* 24, no. 2 (2005): 341-363.

Ruhm, Christopher J. "Recessions, healthy no more?." *Journal of health economics* 42 (2015): 17-28.

Stevens, Ann H., Douglas L. Miller, Marianne E. Page, and Mateusz Filipiński. "The best of times, the worst of times: Understanding pro-cyclical mortality." *American Economic Journal: Economic Policy* 7, no. 4 (2015): 279-311.

Tefft, Nathan. "Insights on unemployment, unemployment insurance, and mental health." *Journal of Health Economics* 30, no. 2 (2011): 258-264.

Tekin, Erdal, Chandler McClellan, and Karen Jean Minyard. *Health and health behaviors during the worst of times: evidence from the Great Recession*. No. w19234. National Bureau of Economic Research, 2013.

Figure 1 Total Unemployment Rate in Each Recession since Postwar






Figure 2 Unemployment Duration since Postwar



Source: US. Bureau of Labor Statistics
research.stlouisfed.org

Figure 3: Impacts of Great Recession by Demographic Group



Note: Each bar corresponds to the impact of a one percentage point increase in the county-level unemployment rate on the probability of currently being employed in the PSID in percentage points.

Table 1: Descriptive Statistics

	Age 25-55			Age 65 and over		
	Obs.	Mean	Std. Dev.	Obs.	Mean	Std. Dev.
Unemployment Rate % (County-level)	43240	6.95	2.75	2514	6.78	2.54
Being Employed (Individual Level)	43280	0.94	0.23	2346	0.69	0.46
<i>Health Behaviors</i>						
Chronic Drinker	24311	0.25	0.43	1060	0.31	0.46
Heavy Drinker	42842	0.27	0.44	2504	0.08	0.27
Current Drinker	42842	0.70	0.46	2504	0.52	0.50
Heavy Smoker	43163	0.15	0.36	2513	0.06	0.23
Current Smoker	42953	0.20	0.40	2417	0.08	0.27
Vigorous Physical Activity	42923	0.71	0.45	2494	0.42	0.49
Light Physical Activity	42846	0.86	0.35	2477	0.73	0.45
Strengthening Activity	36181	0.40	0.49	2042	0.23	0.42
Housework Hours	43019	10.08	8.87	2482	10.92	10.81
<i>Mental Health Outcome</i>						
K6 Index	35739	2.98	3.50	2016	2.17	3.12
<i>Physical Health Outcomes</i>						
Excellent Health	42964	0.62	0.48	2416	0.48	0.50
Poor Health	42964	0.09	0.28	2416	0.20	0.40
Overweight (BMI \geq 25)	41903	0.63	0.48	2368	0.63	0.48
Obese (BMI \geq 30)	41903	0.26	0.44	2368	0.22	0.41
Severely Obese (BMI \geq 35)	41903	0.09	0.29	2368	0.07	0.26
<i>Demographics</i>						
Age	43280	40.88	8.84	2509	71.98	6.41
Sex	43280	0.52	0.50	2518	0.50	0.50
Married	43275	0.67	0.47	2518	0.59	0.49
Never Married	43275	0.16	0.37	2518	0.02	0.15
Widowed	43275	0.01	0.10	2518	0.23	0.42
Divorced	43275	0.13	0.34	2518	0.15	0.36
Less Than High School	41205	0.07	0.26	2293	0.12	0.32
High School Degree	41205	0.32	0.47	2293	0.34	0.47
College Degree	43280	0.63	0.48	2518	0.57	0.49
White	42608	0.80	0.40	2410	0.86	0.34
Black	42608	0.13	0.33	2410	0.07	0.26
Labor Income (2010 dollars)	43280	51457.38	83021.47	2518	24001.30	59351.64

Notes: Data sources are the PSID and LAUS (BLS).

Table 2: Impact of Unemployment on Health and Health Behaviors, Core Results

	Var. Mean	(1)		(2)	
Chronic Drinker	0.25	0.006**	[0.003]	0.008**	[0.003]
Heavy Drinker	0.27	0.002	[0.002]	0.005*	[0.003]
Current Drinker	0.70	0.005**	[0.002]	0.006**	[0.003]
Heavy Smoker	0.15	-0.004**	[0.002]	-0.002	[0.002]
Current Smoker	0.20	-0.002	[0.002]	0.000	[0.002]
Vigorous Physical Activity	0.71	-0.001	[0.003]	0.000	[0.003]
Light Physical Activity	0.86	0.000	[0.002]	0.001	[0.002]
Strengthening Activity	0.40	-0.003	[0.003]	-0.006*	[0.003]
Housework Hours	10.08	0.001	[0.051]	0.019	[0.057]
K6 Index	2.98	0.053**	[0.024]	0.056**	[0.026]
Excellent Health	0.62	-0.009***	[0.003]	-0.009***	[0.003]
Poor Health	0.09	0.008***	[0.002]	0.007***	[0.002]
Overweight	0.63	-0.002	[0.002]	0.001	[0.003]
Obese	0.26	-0.002	[0.002]	-0.001	[0.002]
Severely Obese	0.09	0.000	[0.001]	0.001	[0.002]
Year Fixed Effects			Yes		Yes
County Fixed Effects			Yes		Yes
State-Specific Linear Trends			No		Yes

Notes: Each cell corresponds to a separate regression. Standard errors, which are reported in brackets, are clustered at the county and year. All the regressions control for age, gender, race, marital status, education, and labor income and are weighted using sample weights from the PSID.

* p<0.1, ** p<0.05, *** p<0.001

Table 3: Impact of Unemployment on Health and Health Behaviors, Ages 65 +

	Var. Mean	(1)		(2)	
Chronic Drinker	0.31	0.003	[0.016]	0.000	[0.019]
Heavy Drinker	0.08	0.009	[0.007]	0.007	[0.007]
Current Drinker	0.52	0.017	[0.011]	0.016	[0.013]
Heavy Smoker	0.06	0.001	[0.005]	0.005	[0.006]
Current Smoker	0.08	0.003	[0.006]	0.006	[0.007]
Vigorous Physical Activity	0.42	0.024*	[0.013]	0.021	[0.015]
Light Physical Activity	0.73	0.013	[0.012]	0.014	[0.013]
Strengthening Activity	0.23	0.020*	[0.011]	0.017	[0.013]
Housework Hours	10.92	-0.134	[0.246]	-0.311	[0.290]
K6 Scale	2.17	0.052	[0.095]	0.038	[0.103]
Excellent Health	0.48	0.025**	[0.012]	0.015	[0.014]
Poor Health	0.20	-0.019**	[0.008]	-0.008	[0.009]
Overweight	0.63	-0.013	[0.009]	-0.002	[0.011]
Obese	0.22	0.002	[0.009]	-0.002	[0.010]
Severely Obese	0.07	0.01	[0.007]	0.015*	[0.008]
Year Fixed Effects			Yes		Yes
County Fixed Effects			Yes		Yes
State-Specific Linear Trends			No		Yes

Notes: Per Table 2.

Table 4: Robustness Checks

Outcome	Whole Sample (Individual FE)				Non-mover (County FE)			
Chronic Drinker	0.006	[0.004]	0.007*	[0.004]	0.006*	[0.003]	0.007*	[0.004]
Heavy Drinker	0.004	[0.003]	0.005	[0.003]	0.003	[0.003]	0.006**	[0.003]
Current Drinker	0.002	[0.003]	0.003	[0.003]	0.003	[0.002]	0.006**	[0.003]
Heavy Smoker	-0.003*	[0.002]	-0.002	[0.002]	-0.005***	[0.002]	-0.004**	[0.002]
Current Smoker	-0.002	[0.002]	-0.001	[0.002]	-0.004**	[0.002]	-0.002	[0.002]
Vigorous Physical Activity	0.004	[0.003]	0.004	[0.004]	0.000	[0.003]	0.002	[0.003]
Light Physical Activity	-0.001	[0.003]	0.001	[0.003]	-0.001	[0.003]	0.002	[0.003]
Strengthening Activity	0.000	[0.003]	-0.002	[0.004]	0.001	[0.003]	0.000	[0.003]
Housework Hours	-0.057	[0.058]	-0.043	[0.061]	-0.01	[0.057]	-0.009	[0.064]
K6 Scale	0.028	[0.028]	0.029	[0.030]	0.044*	[0.026]	0.056*	[0.029]
Excellent Health	-0.006*	[0.003]	-0.008**	[0.004]	-0.009***	[0.003]	-0.010***	[0.004]
Poor Health	0.007***	[0.002]	0.008***	[0.002]	0.007***	[0.002]	0.007***	[0.002]
Overweight	0.001	[0.002]	0.003	[0.002]	-0.001	[0.002]	0.001	[0.003]
Obese	-0.003	[0.002]	-0.002	[0.002]	-0.005**	[0.002]	-0.005*	[0.003]
Severely Obese	0.000	[0.002]	0.001	[0.002]	-0.003	[0.002]	-0.002	[0.002]
Year Fixed Effects	Yes		Yes		Yes		Yes	
Type of Fixed Effects	Individual		Individual		County		County	
State-specific Linear Trends	No		Yes		No		Yes	

Notes: Per Table 2.

Table 5: Impact of Unemployment on Health and Health Behaviors by Race

Outcome	White				Black			
Chronic Drinker	0.007**	[0.003]	0.009**	[0.004]	0.012	[0.008]	0.022**	[0.009]
Heavy Drinker	0.000	[0.003]	0.001	[0.003]	0.011*	[0.006]	0.017***	[0.006]
Current Drinker	0.004	[0.002]	0.005*	[0.003]	0.012	[0.007]	0.018**	[0.007]
Heavy Smoker	-0.006***	[0.002]	-0.003	[0.002]	0.006	[0.005]	0.007	[0.006]
Current Smoker	-0.004*	[0.002]	-0.001	[0.002]	0.011**	[0.005]	0.013**	[0.006]
Vigorous Physical Activity	0.001	[0.003]	0.002	[0.003]	0.004	[0.007]	0.005	[0.007]
Light Physical Activity	0.001	[0.002]	0.002	[0.003]	-0.004	[0.006]	-0.005	[0.007]
Strengthening Activity	-0.004	[0.003]	-0.007*	[0.004]	0.004	[0.007]	0.002	[0.007]
Housework Hours	-0.047	[0.051]	-0.001	[0.057]	0.086	[0.112]	-0.054	[0.113]
K6 Index	0.060**	[0.025]	0.068**	[0.028]	-0.048	[0.075]	-0.028	[0.080]
Excellent Health	-0.009***	[0.003]	-0.009**	[0.004]	-0.015*	[0.008]	-0.010	[0.008]
Poor Health	0.008***	[0.002]	0.007***	[0.002]	0.002	[0.005]	-0.001	[0.006]
Overweight	-0.003	[0.003]	0.000	[0.003]	0.014**	[0.005]	0.013**	[0.006]
Obese	-0.005**	[0.002]	-0.003	[0.003]	0.013**	[0.006]	0.011*	[0.007]
Severely Obese	-0.003**	[0.002]	-0.001	[0.002]	0.015***	[0.005]	0.013**	[0.005]
Year Fixed Effects	Yes		Yes		Yes		Yes	
County Fixed Effects	Yes		Yes		Yes		Yes	
State-Specific Linear Trends	No		Yes		No		Yes	

Notes: Per Table 2.

Table 6: Impact of Unemployment on Health and Health Behaviors by Gender

Outcome	Male				Female			
Chronic Drinker	0.005	[0.004]	0.008*	[0.005]	0.008**	[0.004]	0.008**	[0.004]
Heavy Drinker	0.005	[0.003]	0.006	[0.004]	0.001	[0.003]	0.005	[0.004]
Current Drinker	0.007**	[0.003]	0.007**	[0.003]	0.001	[0.003]	0.004	[0.004]
Heavy Smoker	-0.004**	[0.002]	-0.002	[0.002]	-0.003	[0.002]	-0.002	[0.003]
Current Smoker	-0.002	[0.002]	0.000	[0.003]	-0.001	[0.003]	0.000	[0.003]
Vigorous Physical Activity	0.000	[0.003]	0.003	[0.004]	-0.002	[0.004]	-0.003	[0.004]
Light Physical Activity	0.001	[0.003]	0.004	[0.003]	-0.002	[0.003]	0.000	[0.003]
Strengthening Activity	-0.001	[0.004]	-0.005	[0.004]	-0.004	[0.004]	-0.005	[0.005]
Housework Hours	0.047	[0.058]	0.039	[0.065]	-0.035	[0.073]	0.002	[0.078]
K6 Index	0.057**	[0.029]	0.048	[0.031]	0.043	[0.031]	0.051	[0.034]
Excellent Health	-0.007**	[0.004]	-0.007*	[0.004]	-0.009**	[0.004]	-0.009*	[0.004]
Poor Health	0.004*	[0.002]	0.005**	[0.002]	0.010***	[0.003]	0.007**	[0.003]
Overweight	-0.001	[0.003]	0.000	[0.003]	-0.002	[0.003]	0.002	[0.004]
Obese	0.001	[0.003]	0.003	[0.003]	-0.008**	[0.003]	-0.007**	[0.004]
Severely Obese	-0.001	[0.002]	0.001	[0.002]	0.000	[0.002]	0.000	[0.002]
Year Fixed Effects	Yes		Yes		Yes		Yes	
County Fixed Effects	Yes		Yes		Yes		Yes	
State-Specific Linear Trends	No		Yes		No		Yes	

Notes: Per Table 2.

Table 7: Impact of Unemployment on Health and Health Behaviors by Education

Outcome	College				High School			
Chronic Drinker	0.008	[0.005]	0.004	[0.006]	0.006	[0.004]	0.010**	[0.004]
Heavy Drinker	-0.002	[0.004]	-0.001	[0.005]	0.002	[0.003]	0.006	[0.004]
Current Drinker	0.001	[0.004]	0.000	[0.004]	0.006*	[0.003]	0.008**	[0.003]
Heavy Smoker	-0.003	[0.002]	-0.003	[0.002]	-0.003	[0.002]	-0.001	[0.003]
Current Smoker	-0.003	[0.002]	-0.004	[0.003]	-0.001	[0.003]	0.001	[0.003]
Vigorous Physical Activity	-0.001	[0.005]	-0.004	[0.006]	-0.001	[0.004]	0.002	[0.004]
Light Physical Activity	-0.007**	[0.003]	-0.005	[0.004]	0.002	[0.003]	0.004	[0.003]
Strengthening Activity	-0.004	[0.006]	-0.009	[0.006]	-0.003	[0.004]	-0.004	[0.004]
Housework Hours	-0.124	[0.079]	-0.056	[0.086]	0.033	[0.066]	0.01	[0.073]
K6 Index	0.018	[0.037]	0.04	[0.040]	0.064**	[0.030]	0.053	[0.033]
Excellent Health	-0.003	[0.005]	0.000	[0.005]	-0.010***	[0.004]	-0.012***	[0.004]
Poor Health	0.003	[0.002]	0.002	[0.003]	0.008***	[0.002]	0.007***	[0.003]
Overweight	-0.011**	[0.004]	-0.002	[0.005]	0.006*	[0.003]	0.006*	[0.003]
Obese	-0.002	[0.003]	0.000	[0.004]	-0.003	[0.003]	-0.002	[0.003]
Severely Obese	-0.003	[0.002]	-0.002	[0.003]	0.002	[0.002]	0.003	[0.002]
Year Fixed Effects	Yes		Yes		Yes		Yes	
County Fixed Effects	Yes		Yes		Yes		Yes	
State-Specific Linear Trends	No		Yes		No		Yes	

Notes: Per Table 2.